arXiv:hep-ph/0511129v2 3 Feb 2006

MSSM Anatomy of the Polarization Puzzle in $B \to \phi K^*$ Decays

Chao-Shang Huang[a,b], Pyungwon Ko[c], Xiao-Hong Wu[c], Ya-Dong Yang[a]
a Department of Physics, Henan Normal University,
Xinxiang, Henan 453007, China
[b] *Institute of Theoretical Physics,*
Chinese Academy of Science,
P. O. Box 2735, Beijing 100080, China
[c] *School of Physics, Korea Institute for Advanced Study,*
Seoul 130-722, Korea

We analyze the $B \to \phi K^*$ polarization puzzle in the Minimal Supersymmetric Standard Model (MSSM) including the neutral Higgs boson (NHB) contributions. To calculate the non-factorizable contributions to hadronic matrix elements of operators, we have used the QCD factorization framework to the α_s order. It is shown that the recent experimental results of the polarization fractions in $B \to \phi K^*$ decays, which are difficult to be explained in SM, could be explained in MSSM if there are flavor non-diagonal squark mass matrix elements of 2nd and 3rd generations, which also satisfy all relevant constraints from known experiments ($B \to X_s\gamma, B_s \to \mu^+\mu^-, B \to X_s\mu^+\mu^-, B \to X_s g, \Delta M_s$, etc.). We have shown in details that the experimental results can be accommodated with the flavor non-diagonal mass insertion of chirality RL, RL+LR, RR, or LL+ RR when the NHB contributions as well as $\mathcal{O}(\alpha_s)$ corrections of hadronic matrix elements of operators are included. However the branching ratios for the decay are smaller than the experimental measurements.

I. INTRODUCTION

The recent experimental results for polarization fractions in $B \to \phi K^*$ are [1, 2, 3]

$$
\begin{aligned}
|A_0|^2 &= 0.52 \pm 0.08 \pm 0.03 \quad \text{Belle} \\
&= 0.46 \pm 0.12 \pm 0.03 \quad \text{BaBar} \\
|A_\perp|^2 &= 0.19 \pm 0.08 \pm 0.02 \quad \text{Belle}
\end{aligned}
\tag{1}
$$

for the mode ϕK^{*+}, and

$$
\begin{aligned}
|A_0|^2 &= 0.45 \pm 0.05 \pm 0.02 \quad \text{Belle} \\
&= 0.52 \pm 0.05 \pm 0.02 \quad \text{BaBar} \\
|A_\perp|^2 &= 0.30 \pm 0.06 \pm 0.02 \quad \text{Belle} \\
&= 0.22 \pm 0.05 \pm 0.02 \quad \text{BaBar}
\end{aligned}
\tag{2}
$$

for the mode ϕK^{*0}. The amplitudes $|A_0|$ and $|A_{\perp(\|)}|$ are longitudinal and transverse polarizations of decay amplitudes in the transversity basis which satisfy

$$\sum_{i=0,\perp,\|} |A_i|^2 = 1\,. \tag{3}$$

The results deviate significantly from the SM prediction

$$|A_0|^2 \sim 1 - O(1/m_b), \tag{4}$$

based on the naive counting rules which follow from a helicity argument [6]. This significant deviation is referred as a puzzle or anomaly in the literature. It has attracted many interests in searching for possible theoretical explanation in SM and new models beyond SM [6, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 21, 22].

The naive counting rules are obtained with the naive factorization in calculating hadronic matrix elements. It may be possible to explain the data if including the α_s corrections to hadronic matrix elements in SM. It is shown [6] that one can obtain $|A_0|^2 \sim 0.5$ due to the annihilation enhancement from the integral containing end-point singularity in QCDF approach [7]. However, it is at the issue that makes the approach less-predictable. Li and Mishima point out that annihilation contributions are not enough to make $|A_0|^2 \sim 0.5$ [9] in PQCD factorization approach [8]. The effects from the final state interaction (FSI) have been studied in refs. [10, 11]. One can get $|A_0|^2 \sim 0.5$, but $|A_0|^2(B \to \rho K^*) < |A_0|^2(B \to \phi K^*)$ [10] which does not agree with the measurements [1, 4, 5]. Moreover, it has been shown in ref. [11] that such FSI effects would lead to $|A_0|^2 : |A_\||^2 : |A_\perp|^2 = 0.43 : 0.54 : 0.03$ which clearly contradicts the data. Therefore, one may draw the conclusion that it is difficult to explain the data within the SM.

A lot of works have been done to investigate polarizations of $B \to \phi K^*$ in models beyond SM. A model with right currents can give $|A_0|^2 \sim 0.5$ but simultaneously leads to $|A_\||^2 \ll |A_\perp|^2$ which is not in agreement with the data [13, 14]. It is also shown that the RL or LR+RL insertion in MSSM can lead to $|A_0|^2 \sim 0.5$ due to the C_{8g} enhancement, compared with that in SM [15]. However, wrong formulas for the α_s order hadronic matrix elements of the chromomagnetic dipole operator Q_{8g} in the case of transverse polarization are used in Ref. [15]. As shown in refs. [6, 18], the α_s order hadronic matrix elements of Q_{8g} for transverse polarizations are very small. Moreover, the neutral Higgs boson (NHB) contributions are not considered in the work[15]. Yang et al. show that the R-parity violating SUSY might explain the puzzle [20]. A model-independent analysis for contributions of new operators, i.e., the operators beyond the operator basis in SM, has been carried out in ref. [18]. Recently, an analysis of polarizations in the model with scalar interaction of tree-level flavor changing neutral current (e.g., the model III two Higgs doublet model) has also been performed [21]. In this paper, we shall perform a detailed analysis of polarizations in $B \to \phi K^*$ as well as the decay rates in MSSM including neutral Higgs boson contributions and the α_s corrections of hadronic matrix elements.

For the $b \to s$ transition, besides the SM contribution, there are mainly two new contributions arising from the strong penguins and neutral Higgs boson (NHB) penguins with the gluino and squark propagating in the loop in MSSM. The former is not important because the Wilson coefficients of QCD penguin operators in MSSM are not changed significantly, compared with those in SM. Although C_{8g} can get a significant enhancement [23, 27], the hadronic matrix elements of Q_{8g} in the case of transverse polarization are very small. The latter induces scalar operators as well as tensor operators due to renormalization. As well known, the effects of these new operators to leptonic B_s decays are significant [24], and their effects to some hadronic B decays are also important [25, 26]. For $B \to VV$ decays, it is expected that the hadronic matrix elements of scalar and tensor operators can enhance transverse polarization fractions. Moreover, although the effects of the primed counterparts of the usual operators are suppressed by m_s/m_b and consequently negligible in SM, their effects in MSSM can be significant because they have the opposite chirality and the flavor non-diagonal squark mass matrix elements are free parameters which are only subjective to constraints from experiments. In particular, as discussed in ref. [13], the primed counterparts of the usual operators have contributions to longitudinal and transverse polarizations different from those of usual operators and consequently could enhance the transverse polarization fractions. The relevant Wilson coefficients at the m_W scale have been calculated by using the vertex mixing method in Ref.[27] and the mass insertion approximation (MIA) method in ref. [26]. In this paper we shall use the results given in ref. [26]. For the hadronic matrix elements of operators relevant to the decays $B \to VV$, we shall use the BBNS's approach (QCDF) to calculate the α_s order corrections to the naive factorization results.

We show that polarization fractions of the decays can agree with experimental data within 1σ deviation in MSSM with the parameter space satisfying all the constraints from $B_s - \bar{B}_s$ mixing , $B \to X_s\gamma$, $B \to X_s g$, $B \to X_s\mu^+\mu^-$ and $B_s \to \mu^+\mu^-$. In particular, the puzzle for polarization in $B \to \phi K^*$ can be explained, while not in contradiction to the measurements of other two vector final states, in quite a large region of parameter space because we have included the contributions of the primed counterparts of usual operators and NHB induced operators in MSSM with the α_s corrections of their hadronic matrix elements included.

The paper is organized as follows. In Sec. II, we give the effective Hamiltonian responsible for the $b \to s$ transition in MSSM. In Sec. III, we present the decay amplitudes. In particular, the hadronic matrix elements of NHB induced operators to the α_s order are calculated. The Sec. IV is devoted to numerical results. We draw conclusions and discussions in Sec. V.

II. EFFECTIVE HAMILTONIAN

The effective Hamiltonian for $b \to s$ transition can be expressed as[25, 29]

$$\begin{aligned}\mathcal{H}_{\rm eff} &= \frac{G_F}{\sqrt{2}} \sum_{p=u,c} V_{pb} V_{ps}^* \Big(C_1\, Q_1^p + C_2\, Q_2^p + \sum_{i=3,\ldots,16} [C_i\, Q_i + C_i'\, Q_i'] \\ &\quad + C_{7\gamma}\, Q_{7\gamma} + C_{8g}\, Q_{8g} + C_{7\gamma}'\, Q_{7\gamma}' + C_{8g}'\, Q_{8g}' \Big) + \text{h.c.} \end{aligned} \tag{5}$$

Here Q_i are quark and gluon operators and are given by *

$$\begin{aligned}
&Q_1^p = (\bar s_\alpha p_\beta)_{V-A} (\bar p_\beta b_\alpha)_{V-A}, && Q_2^p = (\bar s_\alpha p_\alpha)_{V-A} (\bar p_\beta b_\beta)_{V-A}, \\
&Q_{3(5)} = (\bar s_\alpha b_\alpha)_{V-A} \sum_q (\bar q_\beta q_\beta)_{V-(+)A}, && Q_{4(6)} = (\bar s_\alpha b_\beta)_{V-A} \sum_q (\bar q_\beta q_\alpha)_{V-(+)A}, \\
&Q_{7(9)} = \frac{3}{2} (\bar s_\alpha b_\alpha)_{V-A} \sum_q e_q (\bar q_\beta q_\beta)_{V+(-)A}, && Q_{8(10)} = \frac{3}{2} (\bar s_\alpha b_\beta)_{V-A} \sum_q e_q (\bar q_\beta q_\alpha)_{V+(-)A}, \\
&Q_{11(13)} = (\bar s\, b)_{S+P} \sum_q \frac{m_q}{m_b} (\bar q\, q)_{S-(+)P}\,, \\
&Q_{12(14)} = (\bar s_i\, b_j)_{S+P} \sum_q \frac{m_q}{m_b} (\bar q_j\, q_i)_{S-(+)P}\,, \\
&Q_{15} = \bar s\, \sigma^{\mu\nu} (1+\gamma_5)\, b \sum_q \frac{m_q}{m_b} \bar q\, \sigma_{\mu\nu} (1+\gamma_5)\, q\,,
\end{aligned}$$

---

* For the operators in SM we use the conventions in Ref.[28] where Q_1 and Q_2 are exchanged each other with respect to the convention in most of papers.

$$Q_{16} = \bar{s}_i\,\sigma^{\mu\nu}(1+\gamma_5)\,b_j\sum_q\frac{m_q}{m_b}\bar{q}_j\,\sigma_{\mu\nu}(1+\gamma_5)\,q_i\,,$$
$$Q_{7\gamma} = \frac{e}{8\pi^2}m_b\bar{s}_\alpha\sigma^{\mu\nu}F_{\mu\nu}(1+\gamma_5)b_\beta,$$
$$Q_{8g} = \frac{g_s}{8\pi^2}m_b\bar{s}_\alpha\sigma^{\mu\nu}G^a_{\mu\nu}\frac{\lambda_a^{\alpha\beta}}{2}(1+\gamma_5)b_\beta, \tag{6}$$

where $(\bar{q}_1q_2)_{V\pm A}=\bar{q}_1\gamma^\mu(1\pm\gamma_5)q_2$, $(\bar{q}_1q_2)_{S\pm P}=\bar{q}_1(1\pm\gamma_5)q_2$ †, $p=u,c$, $q=u,d,s,c,b$, e_q is the electric charge number of q quark, λ_a is the color SU(3) Gell-Mann matrix, α and β are color indices, and $F_{\mu\nu}$ ($G_{\mu\nu}$) are the photon (gluon) fields strength. The primed operators, the counterpart of the unprimed operators, are obtained by replacing the chirality in the corresponding unprimed operators with opposite ones.

For the processes we are interested in this paper, the Wilson coefficients should be run down to the scale of $\mathcal{O}(m_b)$. C_1-C_{10} are expanded to $\mathcal{O}(\alpha_s)$ and NLO renormalization group equations (RGEs) should be used. However for the C_{8g} and $C_{7\gamma}$, LO results should be sufficient. The details of the running of these Wilson coefficients can be found in Ref. [29]. The one loop anomalous dimension matrices of the NHB induced operators can be found in refs. [26, 30]. There is the mixing of the new operators induced by NHBs with the operators in SM. The leading order anomalous dimensions have been given in Refs.[31, 32]. The mixing of NHB induced operators with the chromo-magnetic operator can enhance the Wilson coefficient C_{8g} significantly [26, 32]. Because at present no NLO Wilson coefficients $C_i^{(\prime)}$, i=11,...,16, are available, we use the LO running of them in this paper.

III. THE DECAY AMPLITUDE AND POLARIZATION

We use the BBNS approach [7, 28] to calculate the hadronic matrix elements of operators. In the BBNS approach, the hadronic matrix element of an operator in the heavy quark limit can be written as

$$\langle V_1V_2|Q|B\rangle = \langle V_1V_2|Q|B\rangle_f\left[1+\sum r_n\alpha_s^n\right], \tag{7}$$

where $\langle V_1V_2|Q|B\rangle_f$ indicates the naive factorization result. The second term in the square bracket indicates higher order α_s corrections to the matrix elements [28]. We calculate the hadronic matrix elements to the α_s order in this paper. In order to see explicitly the effects of new operators in the MSSM, we divide the decay amplitude into three parts. The first one, H_o, has the same form as that in SM, the second, $H_{o'}$, is for primed counterparts of the SM operators, and the third, H_n, is new which comes from the contributions of Higgs penguin induced operators. That is, we can write the decay amplitude for $B\to VV$ as

$$A(B\to V_1V_2) = \frac{G_F}{\sqrt{2}}H^\lambda$$
$$H^\lambda = H_o^\lambda + H_{o'}^\lambda + H_n^\lambda\,. \tag{8}$$

The helicity amplitudes can be obtained by set $\lambda=0,+1,-1$ in above expressions, respectively.

---

† Strictly speaking, the sum over q in expressions of Q_i (i=11,...,16) should be separated into two parts: one is for q=u, c, i.e., upper type quarks, the other for q=d, s, b, i.e., down type quarks, because the couplings of upper type quarks to NHBs are different from those of down type quarks. In the case of large $\tan\beta$ the former is suppressed by $\tan^{-1}\beta$ with respect to the latter and consequently can be neglected. Hereafter we use, e.g., C_{11}^c to denote the Wilson coefficient of the operator $Q_{11}=(\bar{s}\,b)_{S+P}\,\frac{m_c}{m_b}(\bar{c}\,c)_{S-P}$.

A. Helicity amplitude

Let $e_i^{(\lambda_i)}$ (i=1, 2) be the polarization vector of vector meson V_i, $\lambda_1 = \lambda_2 = \lambda$ in $B \to V_1 V_2$ due to the angular momentum conservation. The helicity amplitudes H^λ $(\lambda = 0, +1, -1)$ of $\bar{B} \to \phi \bar{K^*}^0$ in MSSM are given by

$$\begin{aligned}
H^\lambda &= H_o^\lambda + H_{o'}^\lambda + H_n^\lambda \,, && (9)\\
H_o^\lambda &= -\lambda_t \left[a_3^\lambda + a_4^\lambda + a_5^\lambda - \frac{1}{2}(a_7^\lambda + a_9^\lambda + a_{10}^\lambda)\right] \mathcal{A}_{(V-A)}(\lambda)\,, && (10)\\
H_{o'}^\lambda &= H_o^\lambda \left(C_i \to C_i'\,, \mathcal{A}_{(V-A)}(\lambda) \to \mathcal{A}_{(V+A)}(\lambda)\right)\,, && (11)\\
H_n^\lambda &= -\lambda_t \left\{\left[-\frac{1}{8}(a_{14}^\lambda + r_1 a_{14}'^\lambda) + (a_{15}^\lambda + r_1 a_{15}'^\lambda) + \frac{1}{2}(a_{16}^\lambda + r_1 a_{16}'^\lambda)\right] \mathcal{A}_{T(1+\gamma_5)}(\lambda) - \frac{1}{2}(a_{12}^\lambda + r_2 a_{12}'^\lambda)\mathcal{A}_{(V+A)}(\lambda)\right\} && (12)
\end{aligned}$$

where

$$\begin{aligned}
a_i'^\lambda &= a_i^\lambda (C_i \to C_i')\,,\\
r_1 &= \frac{\mathcal{A}_{T(1-\gamma_5)}}{\mathcal{A}_{T(1+\gamma_5)}}\,, \qquad r_2 = \frac{\mathcal{A}_{(V-A)}}{\mathcal{A}_{(V+A)}}\,,\\
\mathcal{A}_{T(1\pm\gamma_5)}(\lambda) &= \langle \phi(e_2^{(\lambda)}, q)|\bar{s}\sigma_{\mu\nu}(1\pm\gamma_5)s|0\rangle\, \langle K^*(e_1^{(\lambda)}, p_{K^*})|\bar{s}\sigma^{\mu\nu}(1\pm\gamma_5)b|B(p_B)\rangle,\\
\mathcal{A}_{(V\pm A)}(\lambda) &= \langle \phi(e_2^{(\lambda)}, q)|\bar{s}\gamma_\mu(1\pm\gamma_5)s|0\rangle\, \langle K^*(e_1^{(\lambda)}, p_{K^*})|\bar{s}\gamma^\mu(1\pm\gamma_5)b|B(p_B)\rangle. && (13)
\end{aligned}$$

In eq.(10) the coefficients $a_i^\lambda, i = 3, 4, 5, 7, 9, 10$ have been given in ref. [6, 17, 18, 20]. Because there are contradicting results on penguin insertion contributions, especially the C_{8g} effect to transversely polarized amplitudes, we revisit this part and confirm the results in ref. [6, 18]. We calculate coefficients in eq. (12) and results are

$$\begin{aligned}
a_{12}^\lambda &= \frac{m_s}{m_b}\{C_{12} + \frac{C_{11}}{N_c}\} + \frac{\alpha_s}{4\pi}\frac{C_F}{}\frac{C_{11}}{N_c}P_{11}^\lambda\\
a_{14}^\lambda &= \frac{m_s}{m_b}\{C_{14} + \frac{C_{13}}{N_c}[1 + \frac{\alpha_s}{4\pi}\frac{C_F}{}(V_{13}^\lambda + H_{13}^\lambda)]\} + \frac{\alpha_s}{4\pi}\frac{C_F}{}\frac{C_{13}}{N_c}P_{13}^\lambda\\
a_{15}^\lambda &= \frac{m_s}{m_b}\{C_{15} + \frac{C_{16}}{N_c} + \frac{C_{14}}{N_c}\frac{\alpha_s}{4\pi}\frac{C_F}{}(V_{14} + H_{14}^\lambda)\} + \frac{\alpha_s}{4\pi}\frac{C_F}{}\frac{C_{16}}{N_c}P_{16}^\lambda\\
a_{16}^\lambda &= \frac{m_s}{m_b}\{C_{16} + \frac{C_{15}}{N_c}[1 + \frac{\alpha_s}{4\pi}\frac{C_F}{}(V_{15} + H_{15}^\lambda)]\} + \frac{\alpha_s}{4\pi}\frac{C_F}{}\frac{C_{15}}{N_c}P_{15}^\lambda && (14)
\end{aligned}$$

where

$$\begin{aligned}
P_{11}^\lambda &= \left[\frac{m_s}{m_b}(\frac{4}{3}\ln\frac{m_b}{\mu} - G_\phi^\lambda(0)) + \frac{4}{3}\ln\frac{m_b}{\mu} - G_\phi^\lambda(1)\right] r_2,\\
P_{13}^\lambda &= -8\left[-2\ln\frac{m_b}{\mu}\, G_{\phi^\lambda} - GF_\phi^\lambda(1)\right] c_{13}^\lambda,\\
P_{15}^\lambda &= P_{13}^\lambda + 4G_{\phi^\lambda},\\
P_{16}^\lambda &= P_{13}^\lambda, && (15)\\
c_{13}^\lambda &= \begin{cases} r & \text{for } \lambda = 0,\\ \frac{1}{4}\frac{f_\phi}{f_\phi^\perp}\frac{m_\phi}{m_B} & \text{for } \lambda = \pm 1.\end{cases} && (16)
\end{aligned}$$

In eq. (15,16), we have defined

$$\begin{aligned}
&r = \frac{\mathcal{A}_{V-A}(\lambda = 0)}{\mathcal{A}_{T(1+\gamma_5)}(\lambda = 0)}\\
&G_\phi^\lambda(s) = \int_0^1 dx\, G(s - i\epsilon, 1 - x)\, \Phi^\lambda_{\phi\,1}(x)\,, \qquad G(s, x) = -4\int_0^1 dt\, t(1-t)\ln[s - t(1-t)x]\\
&G_{\phi^\lambda} = \int_0^1 \frac{dx}{\bar{x}}\, \Phi^\lambda_{\phi\,2}(x)\,, \quad GF_\phi^\lambda(s) = \int_0^1 dx \frac{\Phi^\lambda_{\phi\,2}(x)}{\bar{x}}\, GF(s - i\,\epsilon, \bar{x})\,, \quad GF(s, x) = \int_0^1 dt \ln[s - x\, t\bar{t}] && (17)
\end{aligned}$$

with $\bar{x} = 1 - x$. Here the distribution amplitudes of ϕ meson are given by

$$\Phi^0_{\phi\,1} = \Phi^0_{\phi\,2} = \phi_\parallel\,, \qquad \Phi^\pm_{\phi\,1} = \left(g_\perp^{(v)} \pm \frac{g_\perp^{\prime(a)}}{4} \right), \qquad \Phi^\pm_{\phi\,2} = \bar{x} \left[g_\perp^{(v)} - \frac{\Phi}{\bar{x}} - \frac{1}{4} \left(\frac{g_\perp^{(a)}}{\bar{x}} \mp g_\perp^{\prime(a)} \right) \right], \tag{18}$$

where $\phi_\parallel\,, g_\perp^{(v)}\,, g_\perp^{(a)}$ and

$$\Phi = \int_0^x dy\, (\phi_\parallel(y) - g_\perp^{(v)}(y)) \tag{19}$$

are defined in ref. [34]. Using the Wandzura-Wilczek approximation [33, 34], one has

$$G_{\phi^\pm} = 0\,. \tag{20}$$

And numerically $GF^+_\phi(1)$ is smaller than $GF^0_\phi(1)$ by about a factor of two and $GF^-_\phi(1) = 0$. Thus, the penguin contract contributions of $Q_{i=13,15,16}$ to transverse amplitudes are smaller than those to the longitudinal amplitude. However, the penguin contract contribution of Q_{11} to transverse amplitudes can be larger than that to the longitudinal amplitude, as $G^+_\phi(1)$ is larger than $G^0_\phi(1)$.

In eq. (14), V^λ and $H^\lambda_{K^*\,\phi}$ are vertex and hard-spectator scattering contributions respectively and numerically not important.

The amplitudes in transversal basis [35] for $\bar{B} \to VV$ are related to the helicity amplitudes by

$$A_0 = H^0\,, \qquad A_\parallel = \frac{H^- + H^+}{\sqrt{2}}\,, \qquad A_\perp = \frac{H^- - H^+}{\sqrt{2}}\,. \tag{21}$$

And the longitudinal polarization is defined by

$$f_L = \frac{|A_0|^2}{|A_0|^2 + |A_\parallel|^2 + |A_\perp|^2}. \tag{22}$$

B. Form factors for $B \to \phi K^*$

Using the identity

$$\sigma^{\mu\nu}\gamma_5 = -\frac{i}{2}\epsilon^{\mu\nu\rho\sigma}\sigma_{\rho\sigma}, \tag{23}$$

we have

$$\begin{aligned} \langle\phi|\bar{s}\sigma^{\mu\nu}(1\pm\gamma_5)s|0\rangle &= (g^{\mu\rho}g^{\nu\sigma} \mp \frac{i}{2}\epsilon^{\mu\nu\rho\sigma})\langle\phi|\bar{s}\sigma_{\rho\sigma}s|0\rangle, \\ \langle K^*(e^{(\lambda)}, p_{K^*})|\bar{s}\sigma^{\mu\nu}(1\pm\gamma_5)b|B(p_B)\rangle &= (g^{\mu\rho}g^{\nu\sigma} \mp \frac{i}{2}\epsilon^{\mu\nu\rho\sigma})\langle K^*(e^{(\lambda)}, p_{K^*})|\bar{s}\sigma_{\rho\sigma}b|B(p_B)\rangle. \end{aligned} \tag{24}$$

Defining

$$\langle K^*(e^{(\lambda)}, p_{K^*})|\bar{s}\sigma_{\mu\nu}b|B(p_B)\rangle = -i\epsilon_{\mu\nu\rho\sigma}\epsilon^{*\rho}(p_B^\sigma C_1(s) + p_{K^*}^\sigma C_2(s)), \tag{25}$$

where $s = q^2$ and $q = p_B - p_{K^*}$, one has

$$\langle K^*(e^{(\lambda)}, p_{K^*})|\bar{s}\sigma^{\mu\nu}q_\nu b|B(p_B)\rangle = i\epsilon^{\mu\nu\rho\sigma}\epsilon_{*\nu}p_{B\,\rho}p_{K^*\,\sigma}(C_1 + C_2). \tag{26}$$

Comparing with the usual definition, one has $(C_1 + C_2) = 2T_1$. From eqs.(24) and (25), it is easy to obtain

$$\begin{aligned} \langle K^*(e^{(\lambda)}, p_{K^*})|\bar{s}\sigma^{\mu\nu}q_\nu(1\pm\gamma_5)b|B(p_B)\rangle = \; & i\epsilon^{\mu\nu\rho\sigma}\epsilon^*_\nu p_{B\,\rho}p_{K^*\,\sigma}C_+ \pm \frac{1}{2}\epsilon^{*\mu}\left[(m_B^2 - m_{K^*}^2)C_+ + sC_-\right] \\ & \mp\frac{1}{2}\epsilon^*\cdot p_B(p^\mu C_+ + q^\mu C_-), \end{aligned} \tag{27}$$

where $C_\pm = C_1 \pm C_2$, $s = q^2$, and $p = p_B + p_{K^*}$. From eq.(27), it follows that there are only two independent form factors in the matrix element of the tensor operator between pseudo-scalar and vector meson states. That is, we need not introduce three form factors in the matrix element, as done in the usual definition in ref. [43]. Comparing with the usual definition of the same matrix element, one has

$$C_+ = 2\,T_1, \;\; C_- = 2\,T_3, \;\; T_2 = \frac{1}{2}\left(C_+ + \frac{s}{m_B^2 - m_{K^*}^2} C_-\right). \tag{28}$$

Define

$$\langle 0|\bar{s}\sigma^{\mu\nu}s|\phi(e^{(\lambda)}, q)\rangle \;=\; f_\phi^\perp (e^{(\lambda)^\mu}\, q^\nu - e^{(\lambda)^\nu}\, q^\mu), \tag{29}$$

we have the naive factorization amplitude of tensor operators as follows.

$$\begin{aligned}\mathcal{A}_{T(1\pm\gamma_5)}(\lambda) \;&\equiv\; \langle\phi(e_2^{(\lambda)},\, q)|\bar{s}\sigma_{\mu\nu}(1\pm\gamma_5)s|0\rangle\langle K^*(e_1^{(\lambda)},\, p_{K^*})|\bar{s}\sigma^{\mu\nu}(1\pm\gamma_5)b|B(p_B)\rangle \\ &=\; 4\,f_\phi^\perp \Big\{ i\,\epsilon_{\mu\nu\rho\sigma} e_2^{*\mu} e_1^{*\nu} p_B^\rho p_{K^*}^\sigma\, 2T_1 \pm (e_1^* \cdot e_2^*)\,(m_B^2 - m_{K^*}^2)\,T_2 \\ &\qquad \mp (e_1^* \cdot p_B)\,(e_2^* \cdot p_B)\, \left(2T_2 + \frac{2q^2}{m_B^2 - m_{K^*}^2} T_3\right) \Big\},\end{aligned} \tag{30}$$

where the superscript (λ) has been suppressed in the right hand of eq. (30). Therefore, it follows that

$$\begin{aligned}\mathcal{A}_{T(1\pm\gamma_5)}(\lambda = 0) \;&=\; \mp 2\,\frac{f_\phi^\perp}{m_\phi m_{K^*}} \Big\{ (m_B^2 - m_\phi^2 - m_{K^*}^2)\,T_2 \;-\; 4m_B^4 p_c^2 \left(T_2 + \frac{m_\phi^2}{m_B^2 - m_{K^*}^2}\,T_3\right) \Big\}, \\ \mathcal{A}_{T(1\pm\gamma_5)}(\lambda = +1) \;&=\; \mp 4\, f_\phi^\perp \;\{ (m_B^2 - m_{K^*}^2)\,T_2 \mp 2m_b p_c\, T_1 \}\,, \\ \mathcal{A}_{T(1\pm\gamma_5)}(\lambda = -1) \;&=\; \mp 4\, f_\phi^\perp \;\{ (m_B^2 - m_{K^*}^2)\,T_2 \pm 2m_b p_c\, T_1 \}\;,\end{aligned} \tag{31}$$

with p_c is the center mass momentum in the $\bar{B}$ rest frame.

The decay constants and the form factors of vector and pseudoscalar mesons are defined as usual [43]:

$$\langle 0|\bar{s}\gamma_\mu(1\pm\gamma_5)s|\phi(q,\, e^{(\lambda)})\rangle \;=\; i\, f_\phi\, m_\phi\, e_\mu^{(\lambda)}\,, \tag{32}$$

$$\begin{aligned}\langle K^*(p_{K^*}, e^{(\lambda)})|\bar{s}\gamma_\mu(1\mp\gamma_5)b|B(p_B)\rangle \;&=\; \epsilon_{\mu\nu\rho\sigma} e^{(\lambda)*\nu} p_B^\rho p_{K^*}^\sigma \frac{2V(s)}{m_B + m_{K^*}} \mp i e_\mu^{(\lambda)*}(m_B + m_{K^*})A_1(s) \\ &\quad \pm i p_\mu e^{(\lambda)*} \cdot p_B \frac{A_2(s)}{m_B + m_{K^*}} \\ &\quad \pm i q_\mu e^{(\lambda)*} \cdot p_B \frac{2m_{K^*}}{s}(A_3(s) - A_0(s))\,,\end{aligned} \tag{33}$$

where $p = p_B + p_{K^*}$ and $q = p_B - p_{K^*}$. The above equations lead to

$$\begin{aligned}\mathcal{A}_{V\pm A}(\lambda) \;&\equiv\; \langle\phi(q,\, e_2^{(\lambda)})|\bar{s}\gamma^\mu(1\pm\gamma_5)s|0\rangle\langle K^*(p_{K^*}, e_1^{(\lambda)})|\bar{s}\gamma_\mu(1\mp\gamma_5)b|B(p_B)\rangle \\ &=\; f_\phi\, m_\phi \left[i\epsilon_{\mu\nu\rho\sigma} e_2^{(\lambda)*\mu} e_1^{(\lambda)*\nu} p_B^\rho p_{K^*}^\sigma \frac{2V(m_\phi^2)}{m_B + m_{K^*}} \pm e_2^{(\lambda)*} \cdot e_1^{(\lambda)*}(m_B + m_{K^*})A_1(m_\phi^2) \right. \\ &\qquad \left. \mp e_1^{(\lambda)*} \cdot p_B e_2^{(\lambda)*} \cdot p_B \frac{A_2(m_\phi^2)}{m_B + m_{K^*}} \right]\end{aligned} \tag{34}$$

¿From eq.(34), we have

$$\begin{aligned}\mathcal{A}_{V\pm A}(\lambda = 0) \;&=\; \mp\frac{f_\phi}{2m_{K^*}} \left\{ (m_B^2 - m_{K^*}^2 - m_\phi^2)(m_B + m_{K^*})\,A_1 - 4m_B^2 p_c^2 \frac{A_2}{m_B + m_{K^*}} \right\}, \\ \mathcal{A}_{V\pm A}(\lambda = +1) \;&=\; \mp f_\phi m_\phi \left\{ (m_B + m_{K^*})\,A_1 \mp 2m_B p_c \frac{V}{m_B + m_{K^*}} \right\}, \\ \mathcal{A}_{V\pm A}(\lambda = -1) \;&=\; \mp f_\phi m_\phi \left\{ (m_B + m_{K^*})\,A_1 \pm 2m_B p_c \frac{V}{m_B + m_{K^*}} \right\}\end{aligned} \tag{35}$$

Comparing eq. (31) and eq. (35), one has

$$\mathcal{A}_{V\mp A} \sim \frac{m_\phi}{m_B}\mathcal{A}_{T(1\pm\gamma_5)}\,. \tag{36}$$

That is, the contributions of tensor operator are enhanced by a factor of m_B/m_ϕ, compared with those of vector-axial vector operators. Therefore, the contributions of NHB are sizable although there is a suppression factor m_s/m_b in eq. (14).

IV. NUMERICAL RESULTS

A. Constraints from experiments

We impose two important constraints from $B \to X_s\gamma$ and $B_s \to \mu^+\mu^-$. Considering the theoretical uncertainties, we take $2.0\times 10^{-4} < \mathrm{Br}(B \to X_s\gamma) < 4.5\times 10^{-4}$, as generally adopted in the literature. Phenomenologically, $\mathrm{Br}(B \to X_s\gamma)$ directly constrains $|C_{7\gamma}(m_b)|^2 + |C'_{7\gamma}(m_b)|^2$ at the leading order. Due to the strong enhancement factor $m_{\tilde g}/m_b$ associated with single $\delta_{23}^{LR(RL)}$ insertion term in $C_{7\gamma}^{(\prime)}(m_b)$, $\delta_{23}^{LR(RL)}$ ($\sim 10^{-2}$) are more severely constrained than $\delta_{23}^{LL(RR)}$. However, if the left-right mixing of scalar bottom quark δ_{33}^{LR} is large (~ 0.5), $\delta_{23}^{LL(RR)}$ is constrained to be order of 10^{-2} since the double insertion term $\delta_{23}^{LL(RR)}\delta_{33}^{LR(LR*)}$ is also enhanced by $m_{\tilde g}/m_b$. The branching ratio $B_s \to \mu^+\mu^-$ in MSSM is given as [24]

$$\begin{aligned}\mathrm{Br}(B_s \to \mu^+\mu^-) \;=\; & \frac{G_F^2\alpha_{\rm em}^2}{64\pi^3} m_{B_s}^3 \tau_{B_s} f_{B_s}^2 |\lambda_t|^2 \sqrt{1-4\widehat{m}^2}[(1-4\widehat{m}^2)|C_{Q_1}(m_b) - C'_{Q_1}(m_b)|^2 + \\ & |C_{Q_2}(m_b) - C'_{Q_2}(m_b) + 2\widehat{m}(C_{10}(m_b) - C'_{10}(m_b))|^2]\end{aligned} \tag{37}$$

where $\widehat{m} = m_\mu/m_{B_s}$. In the moderate and large $\tan\beta$ case the term proportional to $(C_{10} - C'_{10})$ in Eq. (37) can be neglected. The new CDF experimental upper bound of $\mathrm{Br}(B_s \to \mu^+\mu^-)$ is 1.5×10^{-7} [40] at 90% confidence level. We have the constraint

$$\sqrt{|C_{Q_1}(m_W) - C'_{Q_1}(m_W)|^2 + |C_{Q_2}(m_W) - C'_{Q_2}(m_W)|^2} \lesssim 1.2 \tag{38}$$

Because the bound constrains $|C_{Q_i} - C'_{Q_i}|$ (i=1, 2), [‡] we can have values of $|C_{Q_i}|$ and $|C'_{Q_i}|$ larger than those in constrained MSSM (CMSSM) with universal boundary conditions at the high scale and scenarios of the extended minimal flavor violation in MSSM [23] in which $|C'_{Q_i}|$ is much smaller than $|C_{Q_i}|$. Just like the constraint from $\mathrm{Br}(B \to X_s\gamma)$, $\delta_{23}^{LL(RR)}$ is also constrained to be order of 10^{-2} by $\mathrm{Br}(B_s \to \mu^+\mu^-)$, if δ_{33}^{LR} is order of 0.5. At the same time we require that predicted $Br(B \to X_s\mu^+\mu^-)$ falls within 1 σ experimental bounds, which gives no new limits on parameters once the updated CDF bound of $\mathrm{Br}(B_s \to \mu^+\mu^-)$ is imposed. It is shown recently that with the old CDF $\mathrm{Br}(B_s \to \mu^+\mu^-)$ upper bound, 2.6×10^{-6} [42], the present experimental limit R_K ($R_K = Br(B \to K\mu^+\mu^-)/Br(B \to Ke^+e^-)$) ≤ 1.2 puts constraints on $C_{Q_{1,2}}$ which are similar to ones from $\mathrm{Br}(B_s \to \mu^+\mu^-)$ [32] and Higgs penguin contributions (i.e., the terms relevant to $C_{Q_{1,2}}^{(\prime)}$) to $\mathrm{Br}(B \to X_s\mu^+\mu^-)$ is order of 10% or less [32, 41]. We obtained smaller contributions by calculations with the updated CDF bound.

We also impose the current experimental lower bound $\Delta M_s > 14.4\mathrm{ps}^{-1}$ [45]. The correlation between $S_{\phi K}$ and ΔM_s has been extensively discussed in the literature, in particular, in the fourth paper of ref.[23]. So in this paper we just analyze the constraints on parameters from the lower bound. Because $\delta_{23}^{LR(RL)}$ is constrained to be order of 10^{-2} by $\mathrm{Br}(B \to X_s\gamma)$, their contribution to ΔM_s is small. The dominant contribution to ΔM_s comes from $\delta_{23}^{LL(RR)}$ insertion with both constructive and destructive effects compared with the SM contribution. Too large a destructive effect is ruled out, because SM prediction, $\Delta M_s^{SM} = 17.3^{+1.5}_{-0.7}$ [39], is only slightly above the present experiment lower bound. However $\delta_{23}^{LL(RR)}$ are constrained to be order of 10^{-2} by the combined experimental measurement of $\mathrm{Br}(B \to X_s\gamma)$ and upper bound of $\mathrm{Br}(B_s \to \mu^+\mu^-)$. Their effects to ΔM_s are limited. And we have checked, the effects are negligibly small with only one kind of chirality, LL or RR, while ΔM_s can be enhanced to $25\mathrm{ps}^{-1}$ with both kinds of chirality, LL and RR, however it is not strongly correlated with $S_{\phi K_S}$ and $S_{\eta' K_S}$ provided that the other experimental constraints, in particular, those from $\mathrm{Br}(B \to X_s\gamma)$ and upper bound of $\mathrm{Br}(B_s \to \mu^+\mu^-)$, have been imposed.

[‡] $C_{Q_{1,2}}^{(\prime)}$ are the Wilson coefficients of the operators $Q_{1,2}^{(\prime)}$ which are Higgs penguin induced in leptonic and semileptonic B decays and their definition can be found in Ref. [44]. By substituting the quark-Higgs vertex for the lepton-Higgs vertex, it is straightforward to obtain Wilson coefficients relevant to hadronic B decays.

As pointed out in Sec. II, due to the gluino-sbottom loop diagram contribution and the mixing of NHB induced operators with the chromomagnetic dipole operator, the Wilson coefficients $C_{8g}^{(\prime)}$ can be large, which might lead to a too large $Br(B \to X_s g)$. So we need to impose the constraint from experimental upper bound $\mathrm{Br}(B \to X_s g) < 9\%$ [46]. A numerical analysis for C'_{8g}=0 has been performed in Ref.[32]. We carry out a similar analysis by setting both C_{8g} and C'_{8g} non-zero.

B. Numerical results

In the numerical calculations, we employ the latest Light-Cone Sum Rules results [34] for the form factors of $B \to K^*$, other parameters can be found in ref. [26].

Before moving to numerical results, we discuss some unique features of $B \to VV$ process. The contributions of non-primed operators to the helicity amplitude H_+ are much smaller than those to H_-, while the contributions of primed operators to the helicity amplitude H_- are much smaller than those to H_+, because of the helicity flip of quarks and anti-quarks coming from non-primed or primed operators when they consist of a vector meson with some definite helicity. That is, in the transverse basis, A_0 and $A_\parallel$ are proportional to $C - C'$, while $A_\perp$ is proportional to $C + C'$. Therefore, we have $|A_\parallel/A_\perp| \simeq |(C - C')/(C + C')|$.

In numerical analysis we fix $m_{\tilde g} = m_{\tilde q} = 500\mathrm{GeV}$, $\tan\beta = 10$ and $\delta_{33}^{dLR} = 0.4$. We vary the NHB masses in the ranges of $91\mathrm{GeV} \le m_h \le 135\mathrm{GeV}, 91\mathrm{GeV} \le m_H \le 200\mathrm{GeV}$ with $m_h < m_H$ and $200\mathrm{GeV} \le m_A \le 250\mathrm{GeV}$ for the fixed mixing angle $\alpha = 0.6, \pi/2$ of the CP even NHBs and scan δ_{23}^{dAB} in the range $|\delta_{23}^{dAB}| \le 0.06$ for A=B and 0.01 for $A \ne B$ (A = L, R).

Numerical results for the correlation between longitudinal polarization f_L and branching ratio $\mathrm{Br}(B \to \phi K^*)$ are shown in Fig.s 1–4, where the correlation between f_L of $B \to \phi K^*$ and the indirect CP asymmetry $S_{B\to\phi K}$ is also given. Fig. 1, 2, 3, 4 are the results of insertions of δ_{23}^{dRL}, both δ_{23}^{dLR} and δ_{23}^{dRL}, δ_{23}^{dRR}, both δ_{23}^{dLL} and δ_{23}^{dRR}, respectively. In all four cases, f_L can be dragged as low as 0.5, but the $\mathrm{Br}(B \to \phi K^*)$ is smaller than the experimental measurement when $f_L \sim 0.5$. On the other hand, there are some parameter regions with f_L as low as 0.5 and $S_{B\to\phi K}$ near 0.4, which is consistent with the present experimental measurements. In the case of new physics contributions from LR, RL insertions as shown in Fig. 1, 2, the only large effects come from the SUSY contributions of the chromo-magnetic dipole operator Q_{8g} (and/or Q'_{8g}) since the Wilson coefficient $C_{8g}^{\mathrm{new}}(m_b)$ can be significantly larger than $C_{8g}^{\mathrm{SM}}(m_b)$. Because Q_{8g} does not contribute to $h = \pm 1$ amplitude, only the longitudinal amplitude can be largely modified. The experimental measurement of $f_L \sim 0.5$ requires that the magnitude of longitudinal amplitude in MSSM must be smaller than that in SM. Therefore, $\mathrm{Br}(B \to \phi K^*)$ in MSSM decreases, compared with SM, when



FIG. 1: The correlations between f_L and Br$(B \to \phi K^*)$, $S_{B\to\phi K}$ with δ_{23}^{dRL} insertion. The Br is in unit of 10^{-5}.



FIG. 2: The correlations between f_L and Br$(B \to \phi K^*)$, $S_{B\to\phi K}$ with δ_{23}^{dLR} and δ_{23}^{dRL} insertions.

$f_L \sim 0.5$, as can be seen from Fig. 1, 2.

In the case of LL, RR insertions, the Wilson coefficient $C_{8g}^{(\prime)}(m_W)$ could be largely modified and $C_{11}^{(\prime)}(?)$ and $C_{13}^{(\prime)}(?)$ could be large. Running from the m_W scale, $C_{13}(C_{13}^{\prime})$ can induce sizable C_{8g} $(C_{8g}^{\prime})$ at the m_b scale, which effect we have discussed above. Running from a large electro-weak scale to m_b,$C_{13}(C_{13}^{\prime})$ can also induce large C_{13-16} $(C_{13-16}^{\prime})$. However, the updated CDF bound of $B_s \to \mu^+\mu^-$ has imposed a stringent constraint on $C_{11,13}^{(\prime)}$, which leads to that the Wilson coefficients of NHB induced operators are small and $C_{8g}^{(\prime)}$ can be largely modified only for small and moderate $\tan\beta$. The penguin insertions of operators $Q_{13-16}^{(\prime)}$ as well as $Q_{11,12}^{(\prime)}$ have been calculated and given in eqs.(13) and (14), but numerically $Q_{13-16}^{(\prime)}$ contributions to the magnitude of $h = \pm 1$ amplitude are small compared with the magnitude of $h = 0$ amplitude, due to small $GF_{\phi}^{\pm}(1)$ function in eq.(14). $Q_{13-16}^{(\prime)}$ can contribute through tree-level to a_{14-16} in eq.(14) and $Q_{11,12}^{(\prime)}$ also do. However, $C_{11,13}^{(\prime)}$ are not large enough to enhance the transverse amplitudes sizably due to the constraint from $B_s \to \mu^+\mu^-$, as pointed above. So even though $A_\perp$ has the structure of $C + C^{\prime}$, which is different from the $A_{0,\parallel}$ amplitudes, it is still impossible to fine-tune the magnitude of $A_\perp$ to the level of $|A_0|$. Therefore, the overall contribution of LL, RR insertions are very similar to LR, RL insertions, as we see from Fig.s 1–4.

The numerical results are obtained for $m_{\tilde{g}} = m_{\tilde{q}}$=500 GeV. For smaller gluino and squark masses, the Wilson coefficient $C_{8g}^{(\prime)}$ becomes larger, which could have larger effect on the b to s transitions. However, the effect is indeed limited due to the constraint from $B \to X_s g$. For fixed $m_{\tilde{g}}$, the Wilson coefficient $C_{8g}^{(\prime)}$ is not sensitive to the variation of the mass of squark in the range about from 100 GeV to 1.5 TeV. Therefore, the numerical results are not sensitive to the squark mass and would have a sizable change when the gluino mass decreases. When the gluino and squark



FIG. 3: The correlations between f_L and Br($B \to \phi K^*$), $S_{B\to\phi K}$ with δ_{23}^{dRR} insertion.

masses approach to infinity (indeed, the several TeV is big enough), SUSY effects drop, i.e., one reaches the decoupling limit.

Before concluding, we will comment on two channels, $B \to K^*\gamma$ and $B \to K^* l^+ l^-$, which share the same $B \to K^*$ form factors as $B \to K^*\phi$. They have already been calculated within QCD factarization in ref. [50, 51] and new physics effects have been discussed in ref. [52]. For $B \to K^*\gamma$ channel, the SM prediction is about 2 times larger than the experimental measurement. A way out to reduce the theoretical prediction of Br($B \to K^*\gamma$) is to decrease the transverse form factors associated with $B \to K^*$. Then the magnitude of transverse amplitude of $B \to K^*\phi$ will be decreased as well, and the polarization problem becomes even worse within the SM. We carry out an analysis of the correlations between Br($B \to K^*\gamma$) and the polarization of $B \to K^*\phi$ within the new physics framework as we discussed above. We find that both Br($B \to K^*\gamma$) and f_L can be accommodated within 1σ limits only in the case of both LR and RL insertions as shown in Fig. 5a. However, in all the cases, the predicted Br($B \to K^*\phi$) is still small when f_L approaches 0.5. This situation can be relaxed to some extent in all the cases of insertions when we consider the $B \to K^*$ form factors $\xi_{\parallel}$ and $\xi_{\perp}$, as defined in [51], with 50% uncertainties. As an example, our results of the correlations between f_L and Br($B \to K^*\phi$) are given in Fig. 5b in the case of both LL and RR insertions. At the same time, Br($B \to K^*\gamma$) and f_L can be accommodated within 1σ limits in all the cases of insertions. The situation of $B \to K^* l^+ l^-$ is more inconclusive due to the branching ratio measurement by BaBar and Belle with large uncertainties, and theoretically it has been discussed in ref. [51, 52].

V. CONCLUSIONS AND DISCUSSIONS

In summary we have analyzed the $B \to \phi K^*$ polarization puzzle in MSSM. The hadronic matrix elements of the new operators in MSSM for the decays have been calculated in the QCDF approach up to the α_s order. Using the Wilson coefficients in ref. [26] and hadronic matrix elements obtained, we have calculated the polarization fractions and branching ratios for the decays $B \to \phi K^*$. It is shown that in the reasonable region of parameter space where the constraints from $B_s - \bar{B}_s$ mixing , $B \to X_s\gamma$, $B \to X_s g$, $B \to X_s\mu^+\mu^-$ and $B_s \to \mu^+\mu^-$ are satisfied, the polarization fractions of the decays can agree with experimental data within 1σ deviation. In particular, the puzzle for polarization in $B \to \phi K^*$ can be explained, while not in contradiction to the measurements of other two vector final states, in a large region of parameter space because we have included the contributions of the primed operators



FIG. 4: The correlations between f_L and Br$(B \to \phi K^*)$, $S_{B\to\phi K}$ with both δ_{23}^{dLL} and δ_{23}^{dRR} insertions.



a

b

FIG. 5: (a) The correlations between f_L and Br$(B \to K^*\gamma)$ with both δ_{23}^{dLR} and δ_{23}^{dRL} insertions, (b) the correlations between f_L and Br$(B \to K^*\phi)$ with both δ_{23}^{dLL} and δ_{23}^{dRR} insertions, while the $B \to K^*$ form factors $\xi_\parallel$ and $\xi_\perp$ with 50% uncertainties.

and new operators including the α_s corrections of hadronic matrix elements of them in MSSM. However the branching ratio is smaller than the measurements when the longitudinal fraction f_L is near 0.5. We may not worry about it too much at present due to the large uncertainty in calculating hadronic matrix elements of operators.

It is necessary to make a theoretical prediction in SM as precise as we can in order to give a firm ground for signaling new physics. The twist-3 and weak annihilation contributions to $B \to \phi K_S$ in SM have been calculated in Ref. [25] using the method in Ref. [47] by which there is not any phenomenological parameter introduced. The numerical results show that the annihilation contributions to the decay rates are negligible, the twist-3 contributions are also very small, smaller than one percent. We expect that the conclusion would qualitatively remain for $B \to \phi K^*$ in MSSM, so that we have neglected the annihilation contributions in numerical calculations.

In conclusion, we have shown that the recent experimental measurements on the polarization fractions in $B \to \phi K^*$, which is difficult to be explained in SM, can be explained in MSSM if there are flavor non-diagonal squark mass matrix elements of second and third generations whose size satisfies all relevant constraints from known experiments ($B \to X_s\gamma, B_s \to \mu^+\mu^-, B \to X_s\mu^+\mu^-, B \to X_s g, \Delta M_s$, etc.). Therefore, if the present polarization puzzle persists in the future, it will be a signal for new physics beyond the SM and MSSM will be a possible candidate of new physics.

Acknowledgement

The work was supported in part by the Natural Science Foundation of China (NSFC), and in part by KOSEF Sundo grant R02-2003-000-10085-0, and KOSEF through CHEP at Kyungpook National University (PK).

Appendix. Vertex and hard scattering contributions

In the Appendix we give the explicit expressions of vertex corrections and hard scattering contributions at the α_s order for NHB induced operators which are not given in the content.

The hard spectator contributions up to the leading twist in Eq. (14) are given as following:

$$\begin{aligned} H^0_{13,14,15} &= 0, \\ H^\pm_{13} &= 4H^\pm_{14}, \qquad H^\pm_{15} = -12 H^\pm_{14}, \\ H^\pm_{14} &= \frac{4\pi^2}{N_c}(1\mp 1)\,\frac{f_B\, f_\phi^\perp f_{K*}^\perp}{\mathcal{A}_{T(1\pm\gamma_5)}}\int_0^1 \frac{d\xi}{\xi}\Phi_B(\xi)\int_0^1 \frac{du}{\bar{u}}\phi_\perp^\phi(u)\int_0^1 \frac{dv}{\bar{v}}\phi_\perp^{K*}(v), \end{aligned}$$

where $\phi_\perp$ is defined in ref. [34] and normalized as

$$\int_0^1 du\,\phi_\perp(u) = 1.$$

The vertex corrections up to the leading twist in Eq. (14) are as follows.

$$\begin{aligned} V^0_{13,14,15} &= 0 \\ V^\pm_{14} &= -\frac{1}{6}[12\ln\frac{m_b}{\mu} + \int_0^1 du\, g(u)\,\phi_\perp(u)] + the\ regularization\ scheme\ dependent\ constant, \\ g(x) &= 3\left(\frac{1-2x}{1-x}\ln x - i\pi\right), \end{aligned}$$

where we have used that $\phi_\perp(u,\bar{u})$ is symmetric with respect to u, $\bar{u}$. Omitting regularization scheme dependent constants, we have

$$V^\pm_{13} \;=\; 4V^\pm_{14}, \qquad V^\pm_{15} = -12 V^\pm_{14}.$$

We have verified that the μ dependance of $a^\pm_{\text{eff}} = -\frac{1}{8}a^\pm_{14} + a^\pm_{15} + \frac{1}{2}a^\pm_{16}$ in eq.(11) has been cancelled up to the order of α_s. That is, $\frac{d}{d\ln\mu}a^\pm_{\text{eff}} = O(\alpha_s^2)$.

References

---

[1] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **91**, (2003) 171802 [hep-ex/0307026].

[2] Aubert et al. (BABAR Collaboration), Phys. Rev. Lett. **93** (2004) 231804 [hep-ex/0408017].

[3] The Belle Collaboration, K. F. Chen et al, Phys. Rev. Lett. **94** (2005) 221804 [hep-ex/0503013].

[4] B. Aubert *et al.* [BABAR Collaboration], hep-ex/0408093.

[5] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **95** (2005) 141801 [hep-ex/0408102].

[6] A. L. Kagan, Phys. Lett. **B601** (2004) 151 [hep-ph/0405134].

[7] M. Beneke et al., Phys. Rev. Lett. **83**(1999) 1914; Nucl. Phys. **B591**(2000) 313.

[8] H.-n. Li and H. Yu, Phys. Rev. Lett. **74** (1995) 4388; H.-n. Li and T. Yeh, Phys. Rev. **D56** (1997) 1615; Y. Y. Keum, H.-n. Li and A. I. Sanda, Phys. Lett. **B504** (2001) 6; Phys. Rev. **D63** (2001) 054008.

[9] H.-n. Li and S. Mishima, hep-ph/0411146.

[10] P. Colangelo, F.De Fazio, T.N. Pham, Phys. Lett. **B 597**, 291 (2004).

[11] H.Y. Cheng, C.K. Chua, A. Soni, Phys. Rev. **D71** (2005) 014030 [hep-ph/0409317.].

[12] H.-n. Li, hep-ph/0411305.

[13] A. L. kagan, hep-ph/0407076.

[14] E. Alvarez et. al., hep-ph/0410096.

[15] A. .K. Giri and R. Mohanta, hep-ph/0412107.

[16] C. S. Kim, Y.-D. Yang, hep-ph/0412364.

[17] H. -Y. Cheng, K. -C. Yang, Phys. Lett. B **511** (2001) 40.

[18] P. K. Das, K.-C. Yang, Phys. Rev. **D71** (2005) 094002[hep-ph/0412313].

[19] W.-S. Hou, M. Nagashima, hep-ph/0408007.

[20] Y.-D. Yang, R.-M. Wang, G.-R. Lu, Phys. Rev. **D72**(2005) 015009[hep-ph/0411211].

[21] C.-H. Chen, C.-Q. Geng, hep-ph/0504145.

[22] S. Baek, A. Datta, P. Hamel, O. F. Hernandez, D. London, Phys. Rev. **D72** (2005) 094008 [hep-ph/0508149].

[23] G.L. Kane, P. Ko, C. Kolda, J.-H. Park, H. Wang, L.-T.Wang, Phys. Rev. Lett. **90**, 141803 (2003) ; Phys. Rev. **D70** (2004) 035015.

[24] C.-S. Huang and Q.-S. Yan, Phys. Lett. **B442** (1998) 209; C.-S. Huang, W. Liao and Q.-S. Yan, Phys. Rev. **D59** (1999) 011701; S.R. Choudhury and N. Gaur, Phys. Lett. **B451** (1999) 86; K.S. Babu and C. Kolda, Phys. Rev. Lett. **84** (2000) 228; C.-S. Huang et al., Phys. Rev. **D63** (2001) 114021; ibid. **64** (2001) (2001) 059902(E); P.H. Chankowski, L. Slawianowska, P.R. **D63** (2001) 054012; C. Bobeth et al., Phys. Rev. **D64** (2001) 074014; hep-ph/0204225; G. Isidori, A. Retico, JHEP **11** (2001) 001; S. Baek, P. Ko and W. Y. Song, Phys. Rev. Lett. **89** (2002) 271801 ; JHEP **0303**, 054 (2003). For a recent review, see, e.g., C. Kolda, *Tsukuba 2004, Supersymmetry and unification of fundamental interactions* 581, [hep-ph/0409205].

[25] J.-F. Cheng, C.-S. Huang and X.-H. Wu, Phys. Lett. **B 585** (2004) 287 [hep-ph/0306086].

[26] J.-F. Cheng, C.-S. Huang and X.-H. Wu, Nucl. Phys. **B701** (2004) 54.

[27] C.-S. Huang and X.-H. Wu, Nucl. Phys. **B657**(2003) 304.

[28] M. Beneke et al., Nucl. Phys. **B606**(2001) 245.

[29] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**(1996) 1125 [arXiv:hep-ph/9512380].

[30] J.A. Bagger, K.T. Matchev and R.J. Zhang, Phys. Lett. **B412**(1997) 77; M. Ciuchini et al., Nucl. Phys. **B523**(1998) 501; C.-S. Huang and Q.-S. Yan, hep-ph/9906493; A.J. Buras, M. Misiak and J. Urban, Nucl.Phys. **B586** (2000) 397.

[31] F. Borzumati, C. Greub, T. Hurth and D. Wyler, Phys. Rev. **D62**(2000) 075005 .

[32] G. Hiller, F. Krüger, hep-ph/0310219.

[33] S. Wandzura and F. Wilczek, Phys. Lett. B **72** (1977) 195.

[34] P. Ball, V.M. Broaun, Phys. Rev. **D58**, 094016 (1998); P. Ball, R. Zwicky, Phys. Rev. **D71**, 014029 (2005).

[35] I. Dunietz, H.R. Quinn, A. Snyder, W. Toki, H.J. Lipkin, Phys. Rev. **D43**, 2193 (1991).

[36] B.V. Geshkenbein and M.V. Terentev, Phys. Lett. **117**(1982) 243 .

[37] R. Kaiser and H. Leutwyler, Proceedings of Workshop on Nonperturbative Methods in Quantum Field Theory, Adelaide, 1998 [hep-ph/9806336];
R. Kaiser and H. Leutwyler, Eur. Phys. J. C **17** (2000) 623 [hep-ph/0007101].

[38] H.-Y. Cheng, Y.-Y. Keum and K.-C. Yang, Phys.Rev.**D65** (2002) 094023.

[39] M. Ciuchini *et al*, J. High Energy Phys. **07**(2001) 13.

[40] The CDF Collaboration, hep-ex/0508036.

[41] P. H. Chankowski and Ł. Sławianowska, Eur. Phys. J. **C33** (2004) 123.

[42] CDF Collaboration, F. Abe et al., Phys. Rev. **D57** (1998) 3811.

[43] A. Ali, P. Ball, L. T. Handoko, G. Hiller, Phys. Rev. D61 (2000) 074024.

[44] Y.-B. Dai, C.-S. Huang and H.-W. Huang, Phys. Lett. **B390**(1997) 257.

[45] A. Stocchi, Nucl. Phys. Proc. Suppl. **117**(2003) 145 [arXiv:hep-ph/0211245].

[46] A. Kagan, hep-ph/9806266; T.E. Coan et al. (CLEO Collaboration), Phys. Rev. Lett. **80** (1998) 1150.

[47] J.-F. Cheng and C.-S. Huang, Phys. Lett. **B554**(2003) 155 .

[48] R. Harnik, D. T. Larson, H. Murayama, A. Pierce, arXiv:hep-ph/0212180.

[49] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D **58**(1998) 114006 [hep-ph/9802409];
T. Feldmann, Int. J. Mod. Phys. A **15**(2000) 159 [hep-ph/9907491].

[50] S. W. Bosch and G. Buchalla, Nucl. Phys. **B 621** (2002) 459; A. Ali, A. Ya. Parkhomenko, Eur. Phys. J. **C 23** (2002) 89.

[51] M. Beneke, Th. Feldmann, D. Seidel, Nucl. Phys. **B 612** (2001) 25; Eur. Phys. J. **C 41** (2005) 173.

[52] F. Kruger and J. Matias, Phys. Rev. D**71** (2005) 094009.